arXiv:1505.02043v1 [math.OA] 8 May 2015

$\mathbb{Z}_n$-equivariant K-theory

Larry B. Schweitzer

May 7, 2015

Abstract

We construct a sequence of $n-1$ cyclic exact sequences that can be used to compute the K-theory of the $C^\star$-algebra crossed product $A \ltimes \mathbb{Z}_n$.

Contents

1 Introduction **2**

2 Embedding the crossed product as a subalgebra of $M_n(A)$ **3**

3 Exact sequences for the K-theory **7**

4 Example of $PSL_2(\mathbb{R})$ acting on H_2 **11**

5 References **16**

1 Introduction

W. L. Paschke's exact sequence for computing the K-theory of the $C^\star$-algebra crossed product $A \ltimes_\alpha \mathbb{Z}_2$ is

$$\begin{tikzcd}
K_0(A_0) \arrow[r] & K_0(A \ltimes_\alpha \mathbb{Z}_2) \arrow[r] & K_0(A_0/J) \arrow[d] \\
K_1(A_0/J) \arrow[u] & K_1(A \ltimes_\alpha \mathbb{Z}_2) \arrow[l] & K_1(A_0) \arrow[l]
\end{tikzcd}, \tag{1}$$

where α is an action of $\mathbb{Z}_2$ on the $C^\star$-aglebra A, A_0 is the fixed point algebra of α, and J is the ideal of A_0 generated by $\{\, xy \,|\, x, y \in A_1 \,\}$, where $A_1 = \{\, x \in A \,|\, \alpha(x) = -x \,\}$ [Pas, 1985]. See [Ich, 1989] for a generalization of (1) to the case of crossed products by compact groups. We present a generalization of (1) to crossed products by $\mathbb{Z}_n$, where n is any positive integer, $n \in \{2, 3, 4, \dots\}$.

2 Embedding the crossed product as a subalgebra of $M_n(A)$

Let α be an action of $\mathbb{Z}_n$ on a $C^\star$-algebra A. Let ξ be a primitive nth root of unity. For $0 \le k \le n-1$, the kth eigenspace of α is

$$A_k = \{\, x \in A \,|\, \alpha(x) = \xi^k \alpha(x) \,\}. \tag{2}$$

Then $A_k^\star = A_{n-k}$. Note that A_0 is the fixed point algebra of α, and each A_k is an A_0-bimodule. Denote by $A_j A_k$ the closed linear span of $\{\, xy \,|\, x \in A_j, y \in A_k \,\}$. Note that $A_j A_k \subseteq A_{j+k}$ and $A_k A_{n-k}$ is a two-sided ideal in A_0.

Let $P_k \colon A \to A_k$ be the projection

$$P_k(x) = \frac{1}{n} \sum_{i=0}^{n-1} \xi^{-ki} \alpha^i(x), \qquad x \in A. \tag{3}$$

Then $\alpha(P_k(x)) = \xi^k P_k(x)$, P_k is onto A_k, $P_j P_k = \delta_{jk} P_k$, and $\sum_{k=0}^{n-1} P_k = I$.

Elements of the crossed product $A \ltimes_\alpha \mathbb{Z}_n$ are denoted by $\sum_{i=0}^{n-1} a_i \lambda_i$, $a_i \in A$, where $i \mapsto \lambda_i$ a unitary representation of $\mathbb{Z}_n$. Multiplication is given by

$$\bigl(a_i \lambda_i\bigr)\bigl(b_j \lambda_j\bigr) = a\alpha^i(b_j)\lambda_{i+j}. \tag{4}$$

Lemma 1. *The linear map $\theta\colon A \ltimes_\alpha \mathbb{Z}_n \to M_n(A)$ defined by*

$$\theta(a_i\lambda_i) = \begin{pmatrix} P_0(a_i) & \xi^{-i}P_1(a_i) & \dots & \xi^{-i(n-1)}P_{n-1}(a_i) \\ P_{n-1}(a_i) & \xi^{-i}P_0(a_i) & \dots & \xi^{-i(n-1)}P_{n-2}(a_i) \\ \vdots & \vdots & \ddots & \vdots \\ P_1(a_i) & \xi^{-i}P_2(a_i) & \dots & \xi^{-i(n-1)}P_0(a_i) \end{pmatrix} \tag{5}$$

is a $\star$-isomorphism of $A \ltimes_\alpha \mathbb{Z}_n$ with the $C^\star$-subalgebra

$$\begin{pmatrix} A_0 & A_1 & \dots & A_{n-1} \\ A_{n-1} & A_0 & \dots & A_{n-2} \\ \vdots & \vdots & \ddots & \vdots \\ A_1 & A_2 & \dots & A_0 \end{pmatrix} \tag{6}$$

of $M_n(A)$.

Proof: First show θ is an algebra homomorphism. Note $\big(\theta(a_i\lambda_i)\big)_{kl} =$

$\xi^{-il}P_{l-k}(a_i)$, $0 \leq k, l, \leq n-1$. Thus

$$\begin{aligned}
\big(\theta(a_i\lambda_i)\theta(b_j\lambda_j)\big)_{kl} &= \sum_{m=0}^{n-1}\big(\theta(a_i\lambda_i)\big)_{km}\big(\theta(b_j\lambda_j)\big)_{ml} \\
&= \sum_{m=0}^{n-1}\xi^{-im}P_{m-k}(a_i)\xi^{-jl}P_{l-m}(b_j) \\
&= \sum_{m=0}^{n-1}\xi^{-im-kl}\left(\frac{1}{n}\sum_{p=0}^{n-1}\xi^{-(m-k)p}\alpha^p(a_i)\right)\left(\frac{1}{n}\sum_{q=0}^{n-1}\xi^{-(l-m)q}\alpha^q(b_j)\right) \\
&= \frac{1}{n^2}\sum_{m,p,q=0}^{n-1}\xi^{-im-jl-(m-k)p-(l-m)q}\alpha^p(a_i)\alpha^q(b_j) \\
&= \frac{1}{n^2}\sum_{p,q=0}^{n-1}\xi^{-jl+kp-lq}\sum_{m=0}^{n-1}\left(\xi^{m(-i-p+q)}\right)\alpha^p(a_i)\alpha^q(b_j) \\
&= \frac{1}{n^2}\sum_{p,q=0}^{n-1}\xi^{-jl+kp-lq}\left(n\delta_{-i-p+q,0}\right)\alpha^p(a_i)\alpha^q(b_j) \\
&= \frac{1}{n}\sum_{p=0}^{n-1}\xi^{-jl+kp-l(i+p)}\alpha^p(a_i)\alpha^{i+p}(b_j) \qquad -i-p+q=0 \Rightarrow q=i+p \\
&= \xi^{-(i+j)l}\left(\frac{1}{n}\sum_{p=0}^{n-1}\xi^{-(l-k)p}\alpha^p\big(a_i\alpha^i(b_j)\big)\right) \\
&= \xi^{-(i+j)l}P_{l-k}\left(a_i\alpha^i(b_j)\right) \qquad \text{definition of } P_{l-k} \\
&= \theta\left(a_i\alpha^i(b_j)\lambda_{i+j}\right)_{kl} \\
&= \theta\left(\big(a_i\lambda_i\big)\big(b_j\lambda_j\big)\right)_{kl}. \qquad (7)
\end{aligned}$$

It follows that $\theta\left(\big(\sum a_i\lambda_i\big)\big(\sum b_j\lambda_j\big)\right) = \theta\big(\sum a_i\lambda_i\big)\theta\big(\sum b_j\lambda_j\big)$. Similarly, one checks that $\theta\left(\big(\sum a_i\lambda_i\big)^\star\right) = \left(\theta\big(\sum a_i\lambda_i\big)\right)^\star$.

To see that θ is one-to-one, note that an inverse map exists. This is because for $\sum a_j\lambda_j \in A \ltimes_\alpha \mathbb{Z}_n$, we can retrieve each a_j via the formula

$$\frac{1}{n}\sum_{k,l=0}^{n-1} \xi^{li}\big(\theta\big(\sum a_j\lambda_j\big)\big)_{kl} = a_i. \tag{8}$$

The map θ is onto since this inverse map is well defined if we replace $\theta\big(\sum a_j\lambda_j\big)$ by an arbitrary element of the subalgebra (6) of $M_n(A)$. $\square$

Let B_0 denote the subalgebra (6) of Lemma 1. More generally, let B_k be the kth lower right corner of B_0,

$$B_k = \begin{pmatrix} A_0 & A_1 & \dots & A_{n-(k+1)} \\ A_{n-1} & A_0 & \dots & A_{n-(k+2)} \\ \vdots & \vdots & \ddots & \vdots \\ A_{k+1} & A_{k+2} & \dots & A_0 \end{pmatrix}, \tag{9}$$

for $0 \le k \le n-1$, which is $n-k$ by $n-k$. Note $B_{n-1} = A_0$. For $0 \le k \le n-2$, let I_k be the two-sided closed ideal $\sum_{i=1}^{n-1-k} A_iA_{n-i}$ in A_0. Note that for $0 \le k \le n-2$,

$$J_k = \begin{pmatrix} I_k & A_1 & \dots & A_{n-(k+1)} \\ A_{n-1} & A_0 & \dots & A_{n-(k+2)} \\ \vdots & \vdots & \ddots & \vdots \\ A_{k+1} & A_{k+2} & \dots & A_0 \end{pmatrix} \tag{10}$$

is a closed two-sided ideal of the $C^\star$-algebra B_k. Thus for $0 \leq k \leq n-2$, we have $n-1$ exact sequences

$$0 \to J_k \to B_k \to B_k/J_k \cong A_0/I_k \to 0. \tag{11}$$

3 Exact sequences for the K-theory

From the $n-1$ exact sequences (11) of $C^\star$-algebras, we get $n-1$ cyclic exact sequences in K-theory

$$\begin{tikzcd} K_0(J_k) \arrow[r] & K_0(B_k) \arrow[r] & K_0(A_0/I_k) \arrow[d] \\ K_1(A_0/I_k) \arrow[u] & K_1(B_k) \arrow[l] & K_1(J_k) \arrow[l] \end{tikzcd} , \tag{12}$$

for $0 \leq k \leq n-2$.

Lemma 2. *Assume A, I_0, I_1, … I_{n-2} all have strictly positive elements. Then A_0 has a countable approximate unit for A. For each $0 \leq k \leq n-2$, J_k has a countable approximate unit, J_k has a strictly positive element, and B_{k+1} is a full corner of J_k.*

Proof: Let $\{e_n\}_{n=0}^{\infty}$ be an approximate unit for I_k, and $\{f_n\}_{n=0}^{\infty}$ an approximate unit for A in A_0. Then

$$\begin{pmatrix} e_n & 0 & \dots & 0 \\ 0 & f_n & \dots & 0 \\ \vdots & \vdots & \ddots & 0 \\ 0 & 0 & \dots & f_n \end{pmatrix} \tag{13}$$

is an approximate unit for

$$J_k = \begin{pmatrix} I_k & A_1 & \dots & A_{n-(k+1)} \\ A_{n-1} & A_0 & \dots & A_{n-(k+2)} \\ \vdots & \vdots & \ddots & \vdots \\ A_{k+1} & A_{k+2} & \dots & A_0 \end{pmatrix}. \tag{14}$$

It suffices to show that $e_n a \to a$ for all $a \in A_1, \dots, A_{n-(k+1)}$. Note that

$$\|e_n a - a\|^2 = \|(e_n a - a)(e_n a - a)^\star\| = \|e_n a a^\star e_n - e_n a a^\star - a a^\star e_n + a a^\star\|. \tag{15}$$

Now $I_k = A_1 A_{n-1} + \dots A_{(n-1)-k} A_{k+1} = A_1 A_1^\star + \dots A_{(n-1)-k} A_{(n-1)-k}^\star$ so $aa^\star \in I_k$. Thus (15) $\to 0$ as $n \to \infty$.

B_{k+1} is a corner of J_k since

$$\begin{pmatrix} 0 & 0 & \dots & 0 \\ 0 & 1 & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & 1 \end{pmatrix} J_k \begin{pmatrix} 0 & 0 & \dots & 0 \\ 0 & 1 & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & 1 \end{pmatrix} = \begin{pmatrix} 0 & 0 \\ 0 & B_{k+1} \end{pmatrix}. \tag{16}$$

B_{k+1} is full since

$$\begin{align} J_kB_{k+1}J_k &\supseteq \begin{pmatrix} 0 & A_1A_0 & \dots & A_{n-(k+1)}A_0 \\ 0 & A_0^2 & \dots & A_{n-(k+2)}A_0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & A_{k+2}A_0 & \dots & A_0^2 \end{pmatrix} J_k \tag{17} \\ &\supseteq \begin{pmatrix} I_k & A_1 & \dots & A_{n-(k+1)} \\ A_{n-1} & A_0 & \dots & A_{n-(k+2)} \\ \vdots & \vdots & \ddots & \vdots \\ A_{k+1} & A_{k+2} & \dots & A_0 \end{pmatrix} \tag{18} \\ &= J_k. \tag{19} \end{align}$$

Here I used $A_iA_0 = A_i$ for $0 \le i \le n-1$. Let $\{e_\gamma\}$ be an approximate unit for A. Note that P_0e_γ is also an approximate unit for A, since

$$\lim_\gamma (P_0e_\gamma)a = \lim_\gamma \frac{1}{n}\sum \alpha^i(e_\gamma)a = \frac{1}{n}\sum a = a. \tag{20}$$

Thus A_0 contains an approximate unit for A, so $A_iA_0 = A_i$ for each i, $0 \le i \le n-1$. $\square$

By Lemma 2 and Corollary 2.6 of [Bro, 1977], we have

$$J_k \otimes \mathcal{K} \cong B_{k+1} \otimes \mathcal{K}, \tag{21}$$

for $0 \leq k \leq n-2$, where $\mathcal{K}$ is the compact operators on a separable Hilbert space. Thus $K_*(J_k) = K_*(B_{k+1})$ and from our $n-1$ exact sequences (12), we get the $n-1$ exact sequences

$$\begin{tikzcd}
K_0(B_1) \arrow[r] & K_0(A \ltimes_\alpha \mathbb{Z}_n) \arrow[r] & K_0(A_0/I_0) \arrow[d] \\
K_1(A_0/I_0) \arrow[u] & K_1(A \ltimes_\alpha \mathbb{Z}_n) \arrow[l] & K_1(B_1) \arrow[l]
\end{tikzcd} \tag{22}$$

$$\begin{tikzcd}
K_0(B_2) \arrow[r] & K_0(B_1) \arrow[r] & K_0(A_0/I_1) \arrow[d] \\
K_1(A_0/I_0) \arrow[u] & K_1(B_1) \arrow[l] & K_1(B_2) \arrow[l]
\end{tikzcd} \tag{23}$$

$$\vdots$$

$$\begin{tikzcd}
K_0(B_{n-2}) \arrow[r] & K_0(B_{n-3}) \arrow[r] & K_0(A_0/I_{n-3}) \arrow[d] \\
K_1(A_0/I_{n-3}) \arrow[u] & K_1(B_{n-3}) \arrow[l] & K_1(B_{n-2}) \arrow[l]
\end{tikzcd} \tag{24}$$

$$\begin{tikzcd}
K_0(A_0) \arrow[r] & K_0(B_{n-2}) \arrow[r] & K_0(A_0/I_{n-2}) \arrow[d] \\
K_1(A_0/I_{n-2}) \arrow[u] & K_1(B_{n-2}) \arrow[l] & K_1(A_0) \arrow[l]
\end{tikzcd} , \tag{25}$$

where I've identified $B_0 \cong A \ltimes_\alpha \mathbb{Z}_n$ in (22) and $B_{n-1} \cong A_0$ in (25). These exact sequences give a recursive procedure for computing $K_*(A \ltimes_\alpha \mathbb{Z}_n)$ in terms of $K_*(A_0)$, $K_*(A_0/I_{n-2})$, $\ldots$ $K_*(A_0/I_0)$ as follows. Use (25) to compute $K_*(B_{n-2})$ in terms of $K_*(A_0)$ and $K_*(A_0/I_{n-2})$. Then use (24), $K_*(A/I_{n-3})$, and $K_*(B_{n-2})$ to compute $K_*(B_{n-3})$, etc. Finally use (22), $K_*(B_1)$, and $K_*(A_0/I_0)$ to compute $K_*(A \ltimes_\alpha \mathbb{Z}_n)$.

Note that this recursive method reduces to Paschke's exact sequence (1) in the case $n = 2$. When $n = 2$, we are left with the single exact sequence

$$\begin{tikzcd}
K_0(B_1) \arrow[r] & K_0(A \rtimes_\alpha \mathbb{Z}_n) \arrow[r] & K_0(A_0/I_0) \arrow[d] \\
K_1(A_0/I_0) \arrow[u] & K_1(A \rtimes_\alpha \mathbb{Z}_n) \arrow[l] & K_1(B_1) \arrow[l]
\end{tikzcd} , \tag{26}$$

where $B_1 = B_{n-1} = A_0$ and $I_0 = A_1^2$. (Note that $\xi = -1$ in (2), when $n = 2$.)

4 Example of $PSL_2(\mathbb{R})$ acting on H_2

Let H_2 be the hyperbolic plane modelled by the open upper half place of $\mathbb{C}$. Take $A = C_0(H_2)$. The group $PSL_2(\mathbb{R})$ acts on H_2 via

$$z \mapsto \frac{ax + b}{cz + d}. \tag{27}$$

The matrix

$$\begin{pmatrix} 0 & 1 \\ -1 & 1 \end{pmatrix} \tag{28}$$

represents an element of order 3 in $PSL_2(\mathbb{R})$. We compute $K_*(A \rtimes_\alpha \mathbb{Z}_3)$, where

$$\mathbb{Z}_3 = \left\langle \begin{pmatrix} 0 & 1 \\ -1 & 1 \end{pmatrix} \right\rangle \tag{29}$$

and

$$\big(\alpha f\big)(z) = f\Bigg(\begin{pmatrix} 0 & 1 \\ -1 & 1 \end{pmatrix}^{-1} z\Bigg) = f\Bigg(\begin{pmatrix} 1 & -1 \\ 1 & 0 \end{pmatrix} z\Bigg) = f(1 - \frac{1}{z}), \tag{30}$$

$f \in A$.

Computation of $K_*(A_0)$. Let

$$T = \begin{pmatrix} 1 & -1 \\ 1 & 0 \end{pmatrix}. \tag{31}$$

Then the fixed-point algebra is $A_0 = \{\, f \in C_0(H_2) \,|\, f(Tz) = f(z)\,\}$. Find an open set $O \subseteq H_2$ (actually a polygon) such that O, TO, and T^2O are pairwise disjoint and $\overline{O \cup TO \cup T^2O} = H_2$ (see Figure 1 and [Leh, 1966]). The two sides S_1, S_2 of O intersect in T's only fixed point $e^{i\pi/3}$. T maps each point in S_1 to the horizontally adjacent point in S_2, and $A_0 \cong \{\, f \in C_0(O \cup S_1 \cup S_2) \,|\, f(z) = f(Tz), z \in S_1\,\}$. Then A_0 is isomorphic to the algebra of continuous functions on the upper half of the sphere S^2, vanishing at the equator (Figure 2). Hence $A_0 \cong C_0(\mathbb{R}^2)$ and

$$K_*(A_0) = \begin{cases} \mathbb{Z} & * = 0 \\ 0 & * = 1. \end{cases} \tag{32}$$



Figure 1: Drawing of region in the hyperbolic plane. *O* is a fundamental region [Leh, 1966] for the action of T (31).



Figure 2: Deformation of region in hyperbolic plane. Identifying the sides S_1 and S_2 folds the region O into an upper hemisphere.

Computation of $K_*(A_0/I_0)$. Note that since A is commutative, $A_1A_2 = A_2A_1$ and therefore $I_0 = A_1A_2 + A_2A_1 = I_1 = A_1A_2$.

Recall $A_i = \{\, f \in C_0(H_2) \,|\, f(Tz) = \xi^i f(z)\}, i = 0, 1, 2$. If $i \neq 0$, every element of A_i must vanish at any fixed point of T. Thus since functions in A_0 do not necessarily vanish at $e^{i\pi/3}$ (recall that $e^{i\pi/3}$ is T's only fixed point), $A_1A_2 \subsetneq A_0$, falling short of equality at the one point. We have $A_0/I_0 \cong \mathbb{C}$, and therefore

$$K_*(A_0/I_0) = \begin{cases} \mathbb{Z} & * = 0 \\ 0 & * = 1. \end{cases} \tag{33}$$

Use exact sequences to find $K_*(A \ltimes_\alpha \mathbb{Z}_3)$. Now follow the recursive procedure (22)-(25) outlined above to find the K-theory of $A \ltimes_\alpha \mathbb{Z}_3$. Our first exact sequence for $K_*(B_1) = K_*\left(\begin{pmatrix} A_0 & A_1 \\ A_2 & A_0 \end{pmatrix}\right)$ is

$$\begin{tikzcd} K_0(A_0) \arrow[r] & K_0(B_1) \arrow[r] & K_0(A_0/I_0) \arrow[d] \\ K_1(A_0/I_0) \arrow[u] & K_1(B_1) \arrow[l] & K_1(A_0) \arrow[l] \end{tikzcd} \tag{34}$$

or

$$\begin{tikzcd} \mathbb{Z} \arrow[r] & K_0(B_1) \arrow[r] & \mathbb{Z} \arrow[d] \\ 0 \arrow[u] & K_1(B_1) \arrow[l] & 0 \arrow[l] \end{tikzcd}. \tag{35}$$

So $K_0(B_1) = \mathbb{Z} \oplus \mathbb{Z}$, $K_1(B_1) = 0$. Our second exact sequence is

$$\begin{tikzcd} K_0(B_1) \arrow[r] & K_0(A \ltimes_\alpha \mathbb{Z}_3) \arrow[r] & K_0(A_0/I_0) \arrow[d] \\ K_1(A_0/I_0) \arrow[u] & K_1(A \ltimes_\alpha \mathbb{Z}_3) \arrow[l] & K_1(B_1) \arrow[l] \end{tikzcd} \tag{36}$$

or

$$\begin{tikzcd} \mathbb{Z} \oplus \mathbb{Z} \arrow[r] & K_0(A \ltimes_\alpha \mathbb{Z}_3) \arrow[r] & \mathbb{Z} \arrow[d] \\ 0 \arrow[u] & K_1(A \ltimes_\alpha \mathbb{Z}_3) \arrow[l] & 0 \arrow[l] \end{tikzcd}. \tag{37}$$

Thus $K_0(A \ltimes_\alpha \mathbb{Z}_3) = \mathbb{Z}^3$, $K_1(A \ltimes_\alpha \mathbb{Z}_3) = 0$.

5 References

[**Bro, 1977**] L.G. Brown, *Stable isomorphism of hereditary subalgebras of $C^\star$-algebras*, Pacific J. Math. **71(2)** (1977), 335-348.

[**Ich, 1989**] R. Ichihara, *An extension of Paschke's theorem and the Mayer-Vietoris type exact sequences*, Math. Japonica **34(1)** (1989), 27-33.

[**Leh, 1966**] J. Lehner, *A short course in automorphic functions*, Holt, Rinehart, and Winston, New York, 1966.

[**Pas, 1985**] W.L. Paschke, *$\mathbb{Z}_2$-equivariant K-theory*, Lecture Notes in Math. **1132** (1985), 362-373.

Web Page: `http://www.svpal.org/~lsch/Math/indexMath.html`.